

3AL

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
18 August 2005 (18.08.2005)

PCT

(10) International Publication Number
WO 2005/075482 A1

(51) International Patent Classification⁷: C07D 487/08,
471/08, A61K 31/4995, 31/551, A61P 25/00

(21) International Application Number:
PCT/EP2005/050405

(22) International Filing Date: 1 February 2005 (01.02.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
PA 2004 00169 4 February 2004 (04.02.2004) DK
60/541,754 5 February 2004 (05.02.2004) US
60/574,946 28 May 2004 (28.05.2004) US
pa 2004 00839 28 May 2004 (28.05.2004) DK

AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (for all designated States except US): NEUROSEARCH A/S [DK/DK]; 93 Pederstrupvej, DK-2750 Ballerup (DK).

(72) Inventors; and

(75) Inventors/Applicants (for US only): PETERS, Dan [SE/DK]; c/o NeuroSearch A/S, 93 Pederstrupvej, DK-2750 Ballerup (DK). OLSEN, Gunnar, M. [DK/DK]; c/o NeuroSearch A/S, 93 Pederstrupvej, DK-2750 Ballerup (DK). NIELSEN, Elsebet, Østergaard [DK/DK]; c/o NeuroSearch A/S, 93 Pederstrupvej, DK-2750 Ballerup (DK). JØRGENSEN, Tino, Dyhring [DK/DK]; c/o NeuroSearch A/S, 93 Pederstrupvej, DK-2750 Ballerup (DK). AHRING, Philip, K. [DK/DK]; c/o NeuroSearch A/S, 93 Pederstrupvej, DK-2750 Ballerup (DK). TIMMERMANN, Daniel, B. [DK/DK]; c/o NeuroSearch A/S, 93 Pederstrupvej, DK-2750 Ballerup (DK).

(74) Common Representative: NEUROSEARCH A/S; Patent Department, 93 Pederstrupvej, DK-2750 Ballerup (DK).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM,

Declaration under Rule 4.17:


— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: DIAZABICYCLIC ARYL DERIVATIVES AS NICOTINIC ACETYLCHOLINE RECEPTOR LIGANDS



(57) Abstract: This invention relates to novel diazabicyclic aryl derivatives represented by Formula (I); which are found to be cholinergic ligands at the nicotinic acetylcholine receptors. Due to their pharmacological profile the compounds of the invention may be useful for the treatment of diseases or disorders as diverse as those related to the cholinergic system of the central nervous system (CNS), the peripheral nervous

system (PNS), diseases or disorders related to smooth muscle contraction, endocrine diseases or disorders, diseases or disorders related to neuro-degeneration, diseases or disorders related to inflammation, pain, and withdrawal symptoms caused by the termination of abuse of chemical substances.

WO 2005/075482 A1

DIAZABICYCLIC ARYL DERIVATIVES AS NICOTINIC ACETYLCHOLINE RECEPTOR LIGANDS

5 This invention relates to novel diazabicyclic aryl derivatives, which are found to be cholinergic ligands at the nicotinic acetylcholine receptors and modulators of the monoamine receptors and transporters. Due to their pharmacological profile the compounds of the invention may be useful for the treatment of diseases or disorders as diverse as those related to the cholinergic system of the central nervous system
10 (CNS), the peripheral nervous system (PNS), diseases or disorders related to smooth muscle contraction, endocrine diseases or disorders, diseases or disorders related to neuro-degeneration, diseases or disorders related to inflammation, pain, and withdrawal symptoms caused by the termination of abuse of chemical substances.

15

BACKGROUND ART

The endogenous cholinergic neurotransmitter, acetylcholine, exert its biological effect via two types of cholinergic receptors, the muscarinic Acetyl Choline Receptors (mAChR) and the nicotinic Acetyl Choline Receptors (nAChR).

20 As it is well established that muscarinic acetylcholine receptors dominate quantitatively over nicotinic acetylcholine receptors in the brain area important to memory and cognition, and much research aimed at the development of agents for the treatment of memory related disorders have focused on the synthesis of muscarinic acetylcholine receptor modulators.

25 Recently, however, an interest in the development of nAChR modulators has emerged. Several diseases are associated with degeneration of the cholinergic system i.e. senile dementia of the Alzheimer type, vascular dementia and cognitive impairment due to the organic brain damage disease related directly to alcoholism.

WO 00/58311 discloses 1,4-diazabicyclo[3.2.2]nonane-4-carboxylates and
30 carboxamide derivatives useful as inhibitors of the nicotinic $\alpha 7$ receptor subtype. Other 1,4-diazabicyclo[3.2.2]nonane-4-methanone derivatives are not disclosed.

SUMMARY OF THE INVENTION


35 The present invention is devoted to the provision novel modulators of the nicotinic and/or of the monoamine receptors, which modulators are useful for the treatment of diseases or disorders related to the cholinergic receptors, and in particular the nicotinic acetylcholine receptor (nAChR), the serotonin receptor (5-HT₁), the

dopamine receptor (DAR) and the norepinephrine receptor (NER), and of the biogenic amine transporters for serotonin (5-HT), dopamine (DA) and norepinephrine (NE).

Due to their pharmacological profile the compounds of the invention may be useful for the treatment of diseases or disorders as diverse as those related to the cholinergic system of the central nervous system (CNS), the peripheral nervous system (PNS), diseases or disorders related to smooth muscle contraction, endocrine diseases or disorders, diseases or disorders related to neuro-degeneration, diseases or disorders related to inflammation, pain, and withdrawal symptoms caused by the termination of abuse of chemical substances.

The compounds of the invention may also be useful as diagnostic tools or monitoring agents in various diagnostic methods, and in particular for *in vivo* receptor imaging (neuroimaging), and they may be used in labelled or unlabelled form.

In its first aspect the invention provides diazabicyclic aryl derivatives of Formula I



15

any of its enantiomers or any mixture of its enantiomers, an N-oxide, a prodrug, or a pharmaceutically-acceptable addition salt thereof, wherein

n is 1, 2 or 3;

A' and A'', independently of one another, represent an aromatic monocyclic and/or polycyclic, carbocyclic and/or heterocyclic group, optionally substituted one or more times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, carboxy, amino-carbonyl (carbamoyl), sulfamoyl and phenyl; or with another monocyclic or polycyclic, carbocyclic or heterocyclic group; which additional monocyclic or polycyclic, carbocyclic or heterocyclic group may optionally be substituted one or more times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, carboxy, amino-carbonyl (carbamoyl), sulfamoyl and phenyl;

B represents a monocyclic heterocyclic group, optionally substituted one or more times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, oxo, carboxy, amino-carbonyl (carbamoyl), *N*-alkyl-amino-carbonyl

(alkyl-carbamoyl), *N,N*-dialkyl-amino-carbonyl, alkyl-carbonyl-amino, alkyl-carbonyl-amino, sulfamoyl, phenyl or benzyl; or a group of formula -NR'-B', -NR'-(C=V)-B' or -NR'-(C=V)-NR''-B'; wherein R' represents hydrogen, alkyl or a group of formula - (C=V)-NR''-B'; R'' represents hydrogen, alkyl, phenyl or benzyl; V represents O, S or NR'''; wherein R''' represents hydrogen, alkyl or cyano; and B' represents hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, phenyl, benzyl or a monocyclic heterocyclic group; which phenyl, benzyl and heterocyclic groups are optionally substituted one or more times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, oxo, carboxy, amino-carbonyl (carbamoyl), *N*-alkyl-amino-carbonyl (alkyl-carbamoyl), *N,N*-dialkyl-amino-carbonyl, alkyl-carbonyl-amino, amino-carbonyl-amino (ureido), *N*-alkyl-amino-carbonyl-amino (*N*-alkyl-ureido), *N,N*-dialkyl-amino-carbonyl-amino (*N,N*-dialkyl-ureido), sulfamoyl, phenyl and benzyl;

15 and

L represents a single (covalent) bond (i.e. L is absent); or a linking group selected from -CH₂-, -CH₂-CH₂-, -CH=CH-, -C≡C-, -Y-(CH₂)_m-, -(CH₂)_m-Y-, -CONR''''-, -NR''''CO-, -NR''''(SO₂)-, and -(SO₂)NR''''-, wherein Y represents -O-, -S-, -SCH₂-, -SO-, -SO₂-, -NR''''-; R'''' represents hydrogen or alkyl; and m is 0, 1, 2 or 3.

20 In a second aspect the invention provides pharmaceutical compositions comprising a therapeutically effective amount of the diazabicyclic aryl derivative of the invention, or a pharmaceutically-acceptable addition salt thereof, together with at least one pharmaceutically-acceptable carrier or diluent.

Viewed from another aspect the invention relates to the use of the 25 diazabicyclic aryl derivative of the invention, or a pharmaceutically-acceptable addition salt thereof, for the manufacture of pharmaceutical compositions/medicaments for the treatment, prevention or alleviation of a disease or a disorder or a condition of a mammal, including a human, which disease, disorder or condition is responsive to modulation of cholinergic receptors.

30 In yet another aspect the invention provides a method for treatment, prevention or alleviation of diseases, disorders or conditions of a living animal body, including a human, which disorder, disease or condition is responsive to modulation of cholinergic receptors, and which method comprises the step of administering to such a living animal body in need thereof a therapeutically effective amount of the 35 diazabicyclic aryl derivative of the invention.


Other objects of the invention will be apparent to the person skilled in the art from the following detailed description and examples.

DETAILED DISCLOSURE OF THE INVENTION

Diazabicyclic Aryl Derivatives

In its first aspect the invention provides diazabicyclic aryl derivatives of

5 Formula I



any of its enantiomers or any mixture of its enantiomers, an N-oxide, a prodrug, or a pharmaceutically-acceptable addition salt thereof, wherein

n is 1, 2 or 3;

10 A' and A'', independently of one another, represent an aromatic monocyclic and/or polycyclic, carbocyclic and/or heterocyclic group, optionally substituted one or more times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, trihalomethoxy, 15 cyano, nitro, amino, carboxy, amino-carbonyl (carbamoyl), sulfamoyl and phenyl; or with another monocyclic or polycyclic, carbocyclic or heterocyclic group; which additional monocyclic or polycyclic, carbocyclic or heterocyclic group may optionally be substituted one or more times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy, 20 alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, carboxy, amino-carbonyl (carbamoyl), sulfamoyl and phenyl;

B represents a monocyclic heterocyclic group, optionally substituted one or more times with substituents selected from the group consisting of alkyl, cycloalkyl, 25 cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, oxo, carboxy, amino-carbonyl (carbamoyl), N-alkyl-amino-carbonyl (alkyl-carbamoyl), N,N-dialkyl-amino-carbonyl, alkyl-carbonyl-amino, alkyl-carbonyl-amino, sulfamoyl, phenyl or benzyl; or a group of formula -NR'-B', -NR'-(C=V)-B' or 30 -NR'-(C=V)-NR''-B'; wherein R' represents hydrogen, alkyl or a group of formula -(C=V)-NR''-B'; R'' represents hydrogen, alkyl, phenyl or benzyl; V represents O, S or NR'''; wherein R''' represents hydrogen, alkyl or cyano; and B' represents hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, phenyl, benzyl or a monocyclic heterocyclic group; which phenyl, benzyl and heterocyclic groups are optionally 35 substituted one or more times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-

alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, oxo, carboxy, amino-carbonyl (carbamoyl), *N*-alkyl-amino-carbonyl (alkyl-carbamoyl), *N,N*-dialkyl-amino-carbonyl, alkyl-carbonyl-amino, amino-carbonyl-amino (ureido), *N*-alkyl-amino-carbonyl-amino (*N*-alkyl-ureido),
 5 *N,N*-dialkyl-amino-carbonyl-amino (*N,N*-dialkyl-ureido), sulfamoyl, phenyl and benzyl; and

L represents a single (covalent) bond (i.e. L is absent); or a linking group selected from $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, $-\text{CH}=\text{CH}-$, $-\text{C}\equiv\text{C}-$, $-\text{Y}-(\text{CH}_2)_m-$, $-(\text{CH}_2)_m\text{Y}-$, $-\text{CONR}''''-$, $-\text{NR}''''\text{CO}-$, $-\text{NR}''''(\text{SO}_2)-$ and $-(\text{SO}_2)\text{NR}''''-$, wherein Y represents $-\text{O}-$, $-\text{S}-$, $-\text{SCH}_2-$,
 10 $-\text{SO}-$, $-\text{SO}_2-$, $-\text{NR}''''-$; R'''' represents hydrogen or alkyl; and m is 0, 1, 2 or 3.

In one embodiment the diazabicyclic aryl derivative is a compound of Formula I, II, III or IV, as defined herein, wherein n is 1, 2 or 3.

In a more preferred embodiment n is 1 or 2.

In a most preferred embodiment n is 2.

15 In another embodiment the diazabicyclic aryl derivative is a compound of Formula I, II, III or IV, as defined herein, wherein L represents a single (covalent) bond (i.e. L is absent); or a linking group selected from $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, $-\text{CH}=\text{CH}-$, $-\text{C}\equiv\text{C}-$, $-\text{Y}-(\text{CH}_2)_m-$, $-(\text{CH}_2)_m\text{Y}-$, $-\text{CONR}''''-$, $-\text{NR}''''\text{CO}-$, $-\text{NR}''''(\text{SO}_2)-$ and $-(\text{SO}_2)\text{NR}''''-$, wherein Y represents $-\text{O}-$, $-\text{S}-$, $-\text{SCH}_2-$, $-\text{SO}-$, $-\text{SO}_2-$, $-\text{NR}''''-$; R'''' represents hydrogen or alkyl;
 20 and m is 0, 1, 2 or 3.

In a more preferred embodiment L represents a linking group selected from $-\text{CH}=\text{CH}-$, $-\text{C}\equiv\text{C}-$, $-\text{OCH}_2-$, $-\text{CONH}-$, $-\text{NHCO}-$, $-\text{NH}(\text{SO}_2)-$ and $-(\text{SO}_2)\text{NH}-$.

In another preferred embodiment L represents a single (covalent) bond (i.e. L is absent).

25 In a third embodiment the diazabicyclic aryl derivative is a compound of Formula I, II, III or IV, as defined herein, wherein A' represents an aromatic monocyclic or polycyclic, carbocyclic or heterocyclic group, optionally substituted one or more times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy,
 30 cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, carboxy, amino-carbonyl (carbamoyl), sulfamoyl and phenyl; or with another monocyclic or polycyclic, carbocyclic or heterocyclic group; which additional monocyclic or polycyclic, carbocyclic or heterocyclic group may optionally be substituted one or more times with substituents selected from the group consisting of
 35 alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, carboxy, amino-carbonyl (carbamoyl), sulfamoyl and phenyl.

In a more preferred embodiment A' represents an aromatic monocyclic carbocyclic group, optionally substituted one or more times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, carboxy, amino-carbonyl (carbamoyl), sulfamoyl and phenyl.

In an even more preferred embodiment A' represents a phenyl or naphthyl group, optionally substituted one or more times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, carboxy, amino-carbonyl (carbamoyl), sulfamoyl and phenyl.


In a most preferred embodiment A' represents a phen-1,4-diyl or naphth-2,6-diyl group.

In another preferred embodiment A' represents an aromatic monocyclic heterocyclic group, optionally substituted one or more times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, carboxy, amino-carbonyl (carbamoyl), sulfamoyl and phenyl.

In a more preferred embodiment A' represents a furanyl, pyrrolyl, isoxazoly, 1,3,4-oxadiazoly, 1,2,3-oxadiazoly, pyridinyl, pyridinyl, pyridazinyl, indolyl, benzofuranyl, benzothienyl, quinoxaliny or benzimidazoly group.


In an even more preferred embodiment A' represents a furan-2,5-diyl, furan-3,5-diyl, pyrrol-2,5-diyl, isoxazol-3,5-diyl, 1,3,4-oxadiazol-2,5-diyl, 1,2,3-oxadiazol-4,5-diyl, pyridin-2,5-diyl, pyridin-2,4-diyl, pyridazin-3,6-diyl, indol-2,5-diyl, benzofuran-2,5-diyl, benzothien-2,5-diyl, quinoxalin-2,6-diyl or benzimidazol-2,5-diyl group.

In a still more preferred embodiment A' represents a group of formula




wherein X and Y, independently of one another, represent N and/or CR¹, wherein R¹ represents hydrogen, alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, cycloalkoxy, cyanoalkyl, halo, haloalkyl, haloalkoxy, cyano, amino, nitro, aryl, aryloxy, heteroaryl or heteroaryloxy; which aryl, aryloxy, heteroaryl or heteroaryloxy may optionally be substituted one or two times with halo, haloalkyl, haloalkoxy, cyano, amino, nitro and/or a group of the formula -NCOR^{''''}, wherein R^{''''} represents hydrogen or alkyl.

In another preferred embodiment A' represents a group of formula



wherein X and Y, independently of one another, represent N and/or CR¹, wherein R¹ represents hydrogen, alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, cycloalkoxy, cyanoalkyl, halo, haloalkyl, haloalkoxy, cyano, amino, nitro, phenyl or phenyloxy.


In a third preferred embodiment A' represents a group of formula



wherein X represents N or CR², wherein R² represents hydrogen, alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, cycloalkoxy, cyanoalkyl, halo, haloalkyl, haloalkoxy, cyano, amino, nitro, aryl, aryloxy, heteroaryl or heteroaryloxy; which aryl, aryloxy, heteroaryl or heteroaryloxy may optionally be substituted one or two times with halo, haloalkyl, haloalkoxy, cyano, amino, nitro and/or a group of the formula -NCOR^{'''}, wherein R^{'''} represents hydrogen or alkyl; and

In a more preferred embodiment R¹ represents hydrogen, alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, cycloalkoxy, cyanoalkyl, halo, haloalkyl, haloalkoxy, cyano, amino, nitro, aryl, aryloxy, heteroaryl or heteroaryloxy; which aryl, aryloxy, heteroaryl or heteroaryloxy may optionally be substituted one or two times with halo, haloalkyl, haloalkoxy, cyano, amino, nitro and/or a group of the formula -NCOR^{'''}, wherein R^{'''} represents hydrogen or alkyl.

In a fourth preferred embodiment A' represents a group of formula



wherein X represents N or CR², wherein R² represents hydrogen, alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, cycloalkoxy, cyanoalkyl, halo, haloalkyl, haloalkoxy, cyano, amino, nitro, phenyl or phenyloxy; and R¹ represents hydrogen, alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, cycloalkoxy, cyanoalkyl, halo, haloalkyl, haloalkoxy, cyano, amino, nitro, phenyl or phenyloxy.

In a most preferred embodiment A' represents furan-2,5-diyl.

In a fourth embodiment the diazabicyclic aryl derivative is a compound of Formula I, II, III or IV, as defined herein, wherein A'' represents an aromatic monocyclic or polycyclic, carbocyclic or heterocyclic group, optionally substituted one or more times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy,

cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, carboxy, amino-carbonyl (carbamoyl), sulfamoyl and phenyl; or with another monocyclic or polycyclic, carbocyclic or heterocyclic group; which additional monocyclic or polycyclic, carbocyclic or heterocyclic group may optionally be substituted one or more times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, carboxy, amino-carbonyl (carbamoyl), sulfamoyl and phenyl.

10 In a more preferred embodiment A" represents an aromatic monocyclic carbocyclic group, optionally substituted one or more times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, carboxy, amino-carbonyl (carbamoyl), sulfamoyl and phenyl.

15 In an even more preferred embodiment A" represents a phenyl or naphthyl group; which aryl group is optionally substituted one or two times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, carboxy, amino-carbonyl (carbamoyl), sulfamoyl and phenyl.

20 In a most preferred embodiment A" represents a phen-1,3-diyl or phen-1,4-diyl group.

In another preferred embodiment A" represents a furanyl group; which group is optionally substituted one or two times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, carboxy, amino-carbonyl (carbamoyl), sulfamoyl and phenyl.

30 In a more preferred embodiment A" represents a furan-2,5-diyl group; which group is optionally substituted one or two times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, carboxy, amino-carbonyl (carbamoyl), sulfamoyl and phenyl.

35 In a fifth embodiment the diazabicyclic aryl derivative is a compound of Formula I, wherein B represents a monocyclic heterocyclic group, optionally substituted one or more times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-

alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, oxo, carboxy, amino-carbonyl (carbamoyl), *N*-alkyl-amino-carbonyl (alkyl-carbamoyl), *N,N*-dialkyl-amino-carbonyl, alkyl-carbonyl-amino, sulfamoyl, phenyl and benzyl.

5 In a more preferred embodiment B represents a monocyclic heterocyclic group selected from pyrrolidinyl, pyrrolinyl, pyrrolyl, and pyridinyl; which monocyclic heterocyclic group is optionally substituted one or two times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-
10 alkoxy, halo, trihalomethyl, cyano, nitro, amino, oxo, carboxy, amino-carbonyl (carbamoyl), *N*-alkyl-amino-carbonyl (alkyl-carbamoyl), *N,N*-dialkyl-amino-carbonyl, alkyl-carbonyl-amino, sulfamoyl and phenyl.

In an even more preferred embodiment B represents a monocyclic heterocyclic group selected from pyrrolidinyl, pyrrolinyl and pyrrolyl; which monocyclic
15 heterocyclic group is optionally substituted one or two times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, cyano, nitro, amino, oxo, carboxy, amino-carbonyl (carbamoyl), *N*-alkyl-amino-carbonyl (alkyl-carbamoyl), *N,N*-dialkyl-amino-carbonyl,
20 alkyl-carbonyl-amino, sulfamoyl and phenyl.

In a still more preferred embodiment B represents a monocyclic heterocyclic group selected from, pyrrolidinyl, 2-pyrrolinyl (4,5-dihydro-pyrrolyl), 3-pyrrolinyl, (2,5-dihydro-pyrrolyl), pyrrolyl and pyridinyl; which monocyclic heterocyclic group is optionally substituted one or two times with substituents selected from the group
25 consisting of alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, cyano, nitro, amino, oxo, carboxy, carbamoyl (amino-carbonyl), alkyl-carbamoyl (*N*-alkyl-amino-carbonyl), (*N,N*-dialkyl-amino-carbonyl), alkyl-carbonyl-amino, sulfamoyl and phenyl.

30 In a yet more preferred embodiment B represents 3-pyrrolinyl (2,5-dihydro-pyrrolyl) or pyridinyl (pyridin-4-yl); which monocyclic heterocyclic group is optionally substituted one or two times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, cyano,
35 nitro, amino, oxo, carboxy, carbamoyl (amino-carbonyl), alkyl-carbamoyl (*N*-alkyl-amino-carbonyl), (*N,N*-dialkyl-amino-carbonyl), alkyl-carbonyl-amino, sulfamoyl and phenyl.

In a yet still more preferred embodiment B represents 3-pyrrolinyl (2,5-dihydro-pyrrolyl) or pyridinyl (pyridin-4-yl); which monocyclic heterocyclic group is

optionally substituted one or two times with substituents selected from the group consisting of alkyl, hydroxy, alkoxy, halo, trihalomethyl, cyano, nitro, amino and/or oxo.

In a most preferred embodiment the diazabicyclic aryl derivative of the invention is

5 5-Hydroxy-1-{4-[5-(1-oxo-1,4-diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-1,5-dihydro-pyrrol-2-one;


1-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-pyrrolidine-2,5-dione N-oxide; or

(1,4-Diaza-bicyclo[3.2.2]non-4-yl)-[5-(4-pyrrol-1-yl-phenyl)-furan-2-yl]-
10 methanone;

or an enantiomer or a mixture of its enantiomers, or a pharmaceutically-acceptable addition salt thereof.

In a sixth embodiment the diazabicyclic aryl derivative is a compound of Formula I, wherein B represents a group of formula -NR'-B', -NR'-(C=V)-B' or -NR'-(C=V)-NR''-B'; wherein R' represents hydrogen, alkyl or a group of formula -(C=V)-NR''-B'; R'' represents hydrogen, alkyl, phenyl or benzyl; V represents O, S or NR'''; wherein R''' represents hydrogen, alkyl or cyano; and B' represents hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, phenyl, benzyl or a monocyclic heterocyclic group; which phenyl, benzyl and heterocyclic groups are optionally substituted one or
20 more times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, oxo, carboxy, amino-carbonyl (carbamoyl), N-alkyl-amino-carbonyl (alkyl-carbamoyl), N,N-dialkyl-amino-carbonyl, alkyl-carbonyl-amino, amino-carbonyl-amino (ureido), N-alkyl-amino-carbonyl-amino (N-alkyl-ureido), N,N-dialkyl-amino-carbonyl-amino (N,N-dialkyl-ureido), sulfamoyl, phenyl and benzyl.

In a seventh embodiment the diazabicyclic aryl derivative is represented by Formula II



30 any of its enantiomers or any mixture of its enantiomers, or a prodrug, or a pharmaceutically-acceptable addition salt thereof, wherein n, A', A'', L, R' and B' are as defined above.

In a more preferred embodiment L represents a single (covalent) bond (i.e. L is absent); R' represents hydrogen or alkyl; and B' represents hydrogen, alkyl,
35 alkenyl, alkynyl, cycloalkyl, cycloalkenyl, phenyl, benzyl or a monocyclic heterocyclic group; which phenyl, benzyl and heterocyclic groups are optionally substituted one,

two or three times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, oxo, carboxy, amino-carbonyl (carbamoyl), *N*-alkyl-amino-carbonyl
 5 (alkyl-carbamoyl), *N,N*-dialkyl-amino-carbonyl, alkyl-carbonyl-amino, sulfamoyl, phenyl and benzyl.

In an even more preferred embodiment B' represents alkyl, phenyl, benzyl, furanyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, thiadiazolyl, imidazolyl, pyrazolyl, pyridinyl, pyrimidinyl or pyridazinyl; which phenyl, benzyl and heterocyclic groups are
 10 optionally substituted one or two times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, oxo, carboxy, amino-carbonyl (carbamoyl), *N*-alkyl-amino-carbonyl (alkyl-carbamoyl), *N,N*-dialkyl-amino-carbonyl,
 15 alkyl-carbonyl-amino, sulfamoyl, phenyl and benzyl.

In a still more preferred embodiment B' represents furanyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, thiadiazolyl, imidazolyl, pyrazolyl, pyridinyl, pyrimidinyl or pyridazinyl; which heterocyclic group may optionally be substituted one or two times with alkyl, hydroxy-alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, alkoxy-alkyl,
 20 cyanoalkyl, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino and/or phenyl.

In another preferred embodiment B' represents alkyl, phenyl, benzyl or pyridinyl; which phenyl, benzyl and pyridinyl are optionally substituted with hydroxy, alkoxy, halo, trifluoromethyl, cyano, nitro, amino, *N*-alkyl-amino-carbonyl (alkyl-carbamoyl), *N,N*-dialkyl-amino-carbonyl, alkyl-carbonyl-amino, sulfamoyl, phenyl or
 25 benzyl.

In a third preferred embodiment B' represents pyridin-2-yl, pyridin-3-yl, pyridin-4-yl; which pyridinyl may optionally be substituted one or two times with alkyl, hydroxy, alkoxy, halo, trihalomethyl, trihalomethoxy, nitro and/or amino.


In a fourth preferred embodiment *n* is 2; L represents a single (covalent)
 30 bond (i.e. L is absent); A' represents a furanyl, oxazolyl or oxadiazolyl group; A'' represents a phenyl group; R' represents hydrogen or alkyl; and B' represents pyridin-2-yl, pyridin-3-yl, pyridin-4-yl; which pyridinyl may optionally be substituted one or two times with alkyl, hydroxy, alkoxy, halo, trihalomethyl, trihalomethoxy, nitro and/or amino.

35 In a most preferred embodiment the diazabicyclic aryl derivative of the invention is

(1,4-Diaza-bicyclo[3.2.2]non-4-yl)-{5-[4-(3-nitro-pyridin-2-ylamino)-phenyl]-furan-2-yl}-methanone;

or an enantiomer or a mixture of its enantiomers, or a pharmaceutically-acceptable addition salt thereof.

In an eighth embodiment the diazabicyclic aryl derivative is represented by Formula III



5

any of its enantiomers or any mixture of its enantiomers, or a prodrug, or a pharmaceutically-acceptable addition salt thereof, wherein n , A' , A'' , L , R' , V and B' are as defined above.

In a more preferred embodiment L represents a single (covalent) bond (i.e. L is absent); R' represents hydrogen or alkyl; V represents O , S or NR''' ; wherein R''' represents hydrogen, alkyl or cyano; and B' represents hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, phenyl, benzyl or a monocyclic heterocyclic group; which phenyl, benzyl and heterocyclic groups are optionally substituted one, two or three times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, oxo, carboxy, amino-carbonyl (carbamoyl), N -alkyl-amino-carbonyl (alkyl-carbamoyl), N,N -dialkyl-amino-carbonyl, alkyl-carbonyl-amino, sulfamoyl, phenyl and benzyl.

In an even more preferred embodiment V represents O , S or NH ; and B' represents alkyl, alkenyl, cycloalkyl, cycloalkenyl, phenyl or benzyl; which phenyl and benzyl groups are optionally substituted one or two times with alkyl, hydroxy, alkoxy, alkoxy-alkyl, alkoxy-alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, amino-carbonyl (carbamoyl), N -alkyl-amino-carbonyl (alkyl-carbamoyl), N,N -dialkyl-amino-carbonyl and/or alkyl-carbonyl-amino; or furanyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, thiadiazolyl, imidazolyl, pyrazolyl, pyridinyl, pyrimidinyl or pyridazinyl; which heterocyclic group may optionally be substituted one or two times with alkyl, hydroxy-alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, alkoxy-alkyl, cyanoalkyl, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino and/or phenyl.

In another preferred embodiment B' represents a group of formula $-CH_3$, $-CH_2CH_3$, $-CH=CH_2$, $-CH=CH-CH=CH_2$, cyclopenta-1-enyl or cyclopenta-2,4-dienyl, phenyl or benzyl; which phenyl and benzyl may optionally be substituted one or two times with alkyl, hydroxy, alkoxy, halo, trihalomethyl, trihalomethoxy, nitro, amino, amino-carbonyl (amido), N -alkyl-amino-carbonyl (N -alkyl-amido), N,N -dialkyl-amino-carbonyl and/or alkyl-carbonyl-amino.

In a third preferred embodiment B' represents phenyl, benzyl or pyridinyl; which phenyl, benzyl and pyridinyl groups are optionally substituted with halo, trifluoromethyl, cyano, nitro, amino, *N*-alkyl-amino-carbonyl (alkyl-carbamoyl), *N,N*-dialkyl-amino-carbonyl, alkyl-carbonyl-amino or sulfamoyl.

5 In a fourth preferred embodiment n is 2; L represents a single (covalent) bond (i.e. L is absent); A' represents a furanyl, oxazolyl or oxadiazolyl group; A'' represents a phenyl group; R' represents hydrogen or alkyl; V represents O, S or NH; and B' represents a group of formula -CH₃, -CH₂CH₃, -CH=CH₂, -CH=CH-CH=CH₂, cyclopenta-1-enyl cyclopenta-2,4-dienyl, phenyl or benzyl; which phenyl and benzyl
10 may optionally be substituted one or two times with alkyl, hydroxy, alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, amino-carbonyl (amido), *N*-alkyl-amino-carbonyl (*N*-alkyl-amido), *N,N*-dialkyl-amino-carbonyl (*N,N*-dialkyl-amido) and/or alkyl-carbonyl-amino.

In a most preferred embodiment the diazabicyclic aryl derivative of the
15 invention is

N-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-benzamide;

N-{3-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-benzamide;

20 N-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-2-nitro-benzamide;

N-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-4-nitro-benzamide;

N-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-
25 nitro-benzamide;


4-Amino-N-{4-[5-(1,4-diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-benzamide;

3-Amino-N-{4-[5-(1,4-diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-benzamide; or

30 N-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-isonicotinamide;

or an enantiomer or a mixture of its enantiomers, or a pharmaceutically-acceptable addition salt thereof.

In a ninth embodiment the diazabicyclic aryl derivative is represented by
35 Formula IV



any of its enantiomers or any mixture of its enantiomers, or a prodrug, or a pharmaceutically-acceptable addition salt thereof, wherein n, A', A'', L, R', R'', V and B' are as defined above.

In a more preferred embodiment L represents a single (covalent) bond (i.e. L is absent); R' represents hydrogen, alkyl or a group of formula $-(C=V)-NR''-B'$; R'' represents hydrogen, alkyl, phenyl or benzyl; V represents O, S or NR'''; wherein R''' represents hydrogen, alkyl or cyano; and B' represents hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, phenyl, benzyl or a monocyclic heterocyclic group; which phenyl, benzyl and heterocyclic groups are optionally substituted one, two or
 10 three times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, oxo, carboxy, amino-carbonyl (carbamoyl), *N*-alkyl-amino-carbonyl (alkyl-carbamoyl), *N,N*-dialkyl-amino-carbonyl, alkyl-carbonyl-amino, amino-carbonyl-
 15 amino (ureido), *N*-alkyl-amino-carbonyl-amino (*N*-alkyl-ureido), *N,N*-dialkyl-amino-carbonyl-amino (*N,N*-dialkyl-ureido), sulfamoyl, phenyl and benzyl.

In another preferred embodiment V represents O, S or NH; and B' represents alkyl, alkenyl, cycloalkyl, cycloalkenyl, phenyl or benzyl; which phenyl and benzyl groups are optionally substituted one or two times with alkyl, hydroxy, alkoxy,
 20 alkoxy-alkyl, alkoxy-alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, amino-carbonyl (carbamoyl), *N*-alkyl-amino-carbonyl (alkyl-carbamoyl), *N,N*-dialkyl-amino-carbonyl, alkyl-carbonyl-amino, amino-carbonyl-amino (ureido), *N*-alkyl-amino-carbonyl-amino (*N*-alkyl-ureido) and/or *N,N*-dialkyl-amino-carbonyl-amino (*N,N*-dialkyl-ureido); or furanyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, thiadiazolyl, imidazolyl,
 25 pyrazolyl, pyridinyl, pyrimidinyl or pyridazinyl; which heterocyclic group may optionally be substituted one or two times with alkyl, hydroxy-alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, alkoxy-alkyl, cyanoalkyl, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino and/or phenyl.

In a third preferred embodiment B' represents a group of formula $-CH_3$,
 30 $-CH_2CH_3$, $-CH=CH_2$, $-CH=CH-CH=CH_2$, cyclopenta-1-enyl or cyclopenta-2,4-dienyl, phenyl or benzyl; which phenyl and benzyl may optionally be substituted one or two times with alkyl, hydroxy, alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, amino-carbonyl (amido), *N*-alkyl-amino-carbonyl (*N*-alkyl-amido), *N,N*-dialkyl-amino-carbonyl (*N,N*-dialkyl-amido) and/or alkyl-carbonyl-amino.

35 In a fourth preferred embodiment B' represents alkyl, phenyl or benzyl; which phenyl and benzyl groups are optionally substituted one or two times with hydroxy, alkoxy, halo, trifluoromethyl, nitro, amino, alkyl-carbonyl-amino, amino-carbonyl-amino (ureido), *N*-alkyl-amino-carbonyl-amino (*N*-alkyl-ureido) and/or *N,N*-dialkyl-amino-carbonyl-amino (*N,N*-dialkyl-ureido).

In a fifth preferred embodiment n is 2; L represents a single (covalent) bond (i.e. L is absent); A' represents a furanyl, oxazolyl, oxadiazolyl, thiazolyl or pyridazinyl group; A'' represents a phenyl group; and R' represents hydrogen, alkyl or $-(C=O)-NH-B'$; R'' represents hydrogen, alkyl, phenyl or benzyl; V represents O, S or NH; and B' represents a group of formula $-CH_3$, $-CH_2CH_3$, $-CH=CH_2$, $-CH=CH-CH=CH_2$, cyclopenta-1-enyl cyclopenta-2,4-dienyl, phenyl or benzyl; which phenyl and benzyl may optionally be substituted one or two times with alkyl, hydroxy, alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, amino-carbonyl (amido), *N*-alkyl-amino-carbonyl (*N*-alkyl-amido), *N,N*-dialkyl-amino-carbonyl (*N,N*-dialkyl-amido) and/or alkyl-carbonyl-amino.

In a sixth preferred embodiment n is 2; L represents a single (covalent) bond (i.e. L is absent); A' represents a furanyl, oxazolyl, oxadiazolyl, thiazolyl or pyridazinyl group; A'' represents a phenyl group; R' represents hydrogen, alkyl or $-(C=O)-NH-B'$; R'' represents hydrogen, alkyl, phenyl or benzyl; V represents O, S or NH; and B' represents a group of formula $-CH_3$, $-CH_2CH_3$, $-CH=CH_2$, $-CH=CH-CH=CH_2$, cyclopenta-1-enyl cyclopenta-2,4-dienyl, phenyl or benzyl; which phenyl and benzyl may optionally be substituted one or two times with alkyl, hydroxy, alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, amino-carbonyl (amido), *N*-alkyl-amino-carbonyl (*N*-alkyl-amido), *N,N*-dialkyl-amino-carbonyl (*N,N*-dialkyl-amido).

In a seventh preferred embodiment n is 2; L represents a single (covalent) bond (i.e. L is absent); A' represents a furanyl, oxazolyl, oxadiazolyl, thiazolyl or pyridazinyl group; A'' represents a phenyl group; and R' represents hydrogen, alkyl or $-(C=O)-NH-B'$; R'' represents hydrogen, alkyl, phenyl or benzyl; V represents O, S or NH; and B' represents a group of formula $-CH_3$, $-CH_2CH_3$, $-CH=CH_2$, $-CH=CH-CH=CH_2$, cyclopenta-1-enyl cyclopenta-2,4-dienyl, phenyl or benzyl; which phenyl and benzyl may optionally be substituted one or two times with alkyl, hydroxy, alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, amino-carbonyl (amido), *N*-alkyl-amino-carbonyl (*N*-alkyl-amido), *N,N*-dialkyl-amino-carbonyl (*N,N*-dialkyl-amido) and/or alkyl-carbonyl-amino.

In an eighth preferred embodiment B' represents alkyl, phenyl, benzyl or pyridyl; which phenyl, benzyl and pyridyl groups are optionally substituted one or two times with substituents selected from the group consisting of hydroxy, alkoxy, halo, trifluoromethyl, nitro, amino, alkyl-carbonyl-amino, *N*-alkyl-amino-carbonyl-amino (*N*-alkyl-ureido), *N,N*-dialkyl-amino-carbonyl-amino (*N,N*-dialkyl-ureido) and sulfamoyl.

In a most preferred embodiment the diazabicyclic aryl derivative of the invention is

1-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-ethyl-urea;

1-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-phenyl-urea;

1-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-(2-nitrophenyl)-urea;

5 1-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-(2-acetylaminophenyl)-urea;

1-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-(2-aminophenyl)-urea;

10 1-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-(5-chloro-2-methoxyphenyl)-thiourea;

1-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-(5-chloro-2-methoxy-phenyl)-urea;

1-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-benzyl-urea;

15 1-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-1'-benzylaminocarbonyl-3-benzyl-urea;

1-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-(2-chlorophenyl)-urea;

20 1-{3-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-phenyl-urea;

1-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-(2-fluorophenyl)-urea;

1-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-(3-fluorophenyl)-urea;

25 1-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-(2-trifluoromethylphenyl)-urea;

1-{2-[3-[4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl]-ureido)-phenyl}-3-ethyl-urea;

30 1-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-(3-trifluoromethylphenyl)-urea; or

1-{3-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-ethyl-urea;

or an enantiomer or a mixture of its enantiomers, or a pharmaceutically-acceptable addition salt thereof.

35 Any combination of two or more of the embodiments described herein is considered within the scope of the present invention.

Definition of Substituents

In the context of this invention an alkyl group designates a univalent saturated, straight or branched hydrocarbon chain. The hydrocarbon chain preferably contain of from one to eighteen carbon atoms (C₁₋₁₈-alkyl), more preferred of from one
5 to six carbon atoms (C₁₋₆-alkyl; lower alkyl), including pentyl, isopentyl, neopentyl, tertiary pentyl, hexyl and isohexyl. In a preferred embodiment alkyl represents a C₁₋₄-alkyl group, including butyl, isobutyl, secondary butyl, and tertiary butyl. In another preferred embodiment of this invention alkyl represents a C₁₋₃-alkyl group, which may in particular be methyl, ethyl, propyl or isopropyl.

10 In the context of this invention a cycloalkyl group designates a cyclic alkyl group, preferably containing of from three to seven carbon atoms (C₃₋₇-cycloalkyl), including cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl.

In the context of this invention a cycloalkyl-alkyl group designates a cycloalkyl group as defined above, which cycloalkyl group is substituted on an alkyl
15 group as also defined above. Examples of preferred cycloalkyl-alkyl groups of the invention include cyclopropylmethyl and cyclopropylethyl.

In the context of this invention an alkoxy group designates an "alkyl-O-" group, wherein alkyl is as defined above. Examples of preferred alkoxy groups of the invention include methoxy and ethoxy.

20 In the context of this invention a cycloalkoxy group designates a "cycloalkyl-O-" group, wherein cycloalkyl is as defined above.

In the context of this invention, a cyano-alkyl group designates an alkyl group substituted with CN, wherein alkyl is as defined above.

In the context of this invention halo represents fluoro, chloro, bromo or
25 iodo, and haloalkyl group designates an alkyl group as defined herein, which alkyl group is substituted one or more times with halo. Thus a trihalomethyl group represents e.g. a trifluoromethyl group, a trichloromethyl group, and similar trihalo-substituted methyl groups. Preferred haloalkyl groups of the invention include trihalogenmethyl, preferably -CF₃.

30 In the context of this invention a haloalkoxy group designates an alkoxy group as defined herein, which alkoxy group is substituted one or more times with halo. Preferred haloalkoxy groups of the invention include trihalogenmethoxy, preferably -OCF₃.

In the context of this invention an aryl group designates a monocyclic or
35 polycyclic aromatic hydrocarbon group. Examples of preferred aryl groups of the invention include phenyl, indenyl, naphthyl, azulenyl, fluorenyl, and anthracenyl. The most preferred aryl group of the invention is phenyl.

In the context of this invention an aryloxy group designates an "aryl-O-" group, wherein aryl is as defined above. The most preferred aryloxy group of the invention is phenoxy.

In the context of this invention a heteroaryl group designates an aromatic
5 mono- or polycyclic heterocyclic group, which holds one or more heteroatoms in its ring structure. Preferred heteroatoms include nitrogen (N), oxygen (O) and sulphur (S).

Preferred 5-6 membered heteroaryl groups of the invention include furanyl, in particular furan-2- or 3-yl; thienyl, in particular thien-2- or 3-yl; selenophenyl, in particular selenophen-2- or 3-yl; pyrrolyl (azolyl), in particular pyrrol-2- or 3-yl;
10 oxazolyl, in particular oxazol-2, 4- or 5-yl; thiazolyl, in particular thiazol-2, 4- or 5-yl; imidazolyl, in particular imidazol-2- or 4-yl; pyrazolyl, in particular pyrazol-3- or 4-yl; isoxazolyl, in particular isoxazol-3, 4- or 5-yl; isothiazolyl, in particular isothiazol-3-, 4- or 5-yl; oxadiazolyl, in particular 1,2,3-oxadiazol-4- or 5-yl, or 1,3,4-oxadiazol-2-yl;
15 triazolyl, in particular 1,2,3-triazol-4-yl or 1,2,4-triazol-3-yl; thiadiazolyl, in particular 1,2,3-thiadiazol-4- or 5-yl, or 1,3,4-thiadiazol-2-yl; pyridyl, in particular pyrid-2-, 3- or 4-yl; pyridazinyl, in particular pyridazin-3- or 4-yl; pyrimidinyl, in particular pyrimidin-2-, 4- or 5-yl; pyrazinyl, in particular pyrazin-2- or 3-yl; and triazinyl, in particular 1,2,4- or 1,3,5-triazinyl.

20 More preferred 5 membered heteroaryl groups of the invention include furanyl, in particular furan-2- or 3-yl; thienyl, in particular thien-2- or 3-yl; pyrrolyl (azolyl), in particular pyrrol-2- or 3-yl; oxazolyl, in particular oxazol-2, 4- or 5-yl; thiazolyl, in particular thiazol-2, 4- or 5-yl; isoxazolyl, in particular isoxazol-3, 4- or 5-yl; isothiazolyl, in particular isothiazol-3-, 4- or 5-yl; and thiadiazolyl, in particular
25 1,2,3-thiadiazol-4- or 5-yl, or 1,3,4-thiadiazol-2-yl.

Yet more preferred 5 membered heteroaryl groups of the invention include furanyl, thienyl, pyrrolyl, oxazolyl and oxadiazolyl.

Most preferred 5 membered heteroaryl groups of the invention include furanyl, in particular furan-2- or 3-yl; and thienyl, in particular thien-2- or 3-yl.

30 More preferred 6 membered heteroaryl groups of the invention include pyridyl, in particular pyrid-2-, 3- or 4-yl; and pyrazinyl, in particular pyrazin-2- or 3-yl.

In the context of this invention an aromatic bicyclic heterocyclic group designates a bicyclic heterocyclic group, which holds one or more heteroatoms in its ring structure. In the context of this invention the term "bicyclic heterocyclic group"
35 includes benzo-fused five- and six-membered heterocyclic rings containing one or more heteroatoms. Preferred heteroatoms include nitrogen (N), oxygen (O), and sulphur (S).

Preferred bicyclic heteroaryl groups of the invention include indolizinyl, in particular indolizin-2-, 5- or 6-yl; indolyl, in particular indol-2-, 5- or 6-yl; isoindolyl, in

particular isoindol-2-, 5- or 6-yl; benzo[b]furanyl, in particular benzofuran-2-, 5- or 6-yl; benzo[b]thienyl, in particular benzothien-2-, 5- or 6-yl; benzoimidazolyl, in particular benzoimidazol-2-, 5- or 6-yl; benzothiazolyl, in particular benzothiazol-5- or 6-yl; purinyl, in particular purin-2- or 8-yl; quinolinyl, in particular quinolin-2-, 3-, 6- or 7-yl; isoquinolinyl, in particular isoquinolin-3-, 6- or 7-yl; cinnolinyl, in particular cinnolin-6- or 7-yl; phthalazinyl, in particular phthalazin-6- or 7-yl; quinazolinyl, in particular quinazolin-2-, 6- or 7-yl; quinoxalinyl, in particular quinoxalin-2- or 6-yl; 1,8-naphthyridinyl, in particular 1,8-naphthyridin-2-, 3-, 6- or 7-yl; and pteridinyl, in particular pteridin-2-, 6- or 7-yl.

10 More preferred bicyclic heteroaryl groups of the invention include indolyl, in particular indol-2-, 5- or 6-yl; benzo[b]furanyl, in particular benzofuran-2-, 5- or 6-yl; benzo[b]thienyl, in particular benzothien-2-, 5- or 6-yl; benzoimidazolyl, in particular benzoimidazol-2-, 5- or 6-yl; and quinoxalinyl, in particular quinoxalin-2- or 6-yl.

Most preferred bicyclic heteroaryl groups of the invention include indolyl, in particular indol-2-, 5- or 6-yl; benzo[b]furanyl, in particular benzofuran-2-, 5- or 6-yl; benzo[b]thienyl, in particular benzothien-2-, 5- or 6-yl.

In the context of this invention a heteroaryloxy group designates a "heteroaryl-O-" group, wherein heteroaryl is as defined above.

20 Pharmaceutically Acceptable Salts


The diazabicyclic aryl derivative of the invention may be provided in any form suitable for the intended administration. Suitable forms include pharmaceutically (i.e. physiologically) acceptable salts, and pre- or prodrug forms of the chemical compound of the invention.

25 Examples of pharmaceutically acceptable addition salts include, without limitation, the non-toxic inorganic and organic acid addition salts such as the hydrochloride, the hydrobromide, the nitrate, the perchlorate, the phosphate, the sulphate, the formate, the acetate, the aconate, the ascorbate, the benzenesulphonate, the benzoate, the cinnamate, the citrate, the embonate, the enantate, the fumarate, the glutamate, the glycolate, the lactate, the maleate, the malonate, the mandelate, the methanesulphonate, the naphthalene-2-sulphonate derived, the phthalate, the salicylate, the sorbate, the stearate, the succinate, the tartrate, the toluene-p-sulphonate, and the like. Such salts may be formed by procedures well known and described in the art.

35 Metal salts of a chemical compound of the invention include alkali metal salts, such as the sodium salt of a chemical compound of the invention containing a carboxy group.

In the context of this invention the "onium salts" of N-containing compounds may also be contemplated as pharmaceutically acceptable salts. Preferred "onium

salts" include the alkyl-onium salts, the cycloalkyl-onium salts, and the cycloalkylalkyl-onium salts. Particularly preferred onium salts of the invention include those created at the N' position according to the following Formula I'



5

Steric Isomers

The chemical compounds of the present invention may exist in (+) and (-) forms as well as in racemic forms. The racemates of these isomers and the individual isomers themselves are within the scope of the present invention.

10 Racemic forms can be resolved into the optical antipodes by known methods and techniques. One way of separating the diastereomeric salts is by use of an optically active acid, and liberating the optically active amine compound by treatment with a base. Another method for resolving racemates into the optical antipodes is based upon chromatography on an optical active matrix. Racemic
15 compounds of the present invention can thus be resolved into their optical antipodes, e.g., by fractional crystallisation of d- or l- (tartrates, mandelates, or camphorsulphonate) salts for example.

The chemical compounds of the present invention may also be resolved by the formation of diastereomeric amides by reaction of the chemical compounds of the
20 present invention with an optically active activated carboxylic acid such as that derived from (+) or (-) phenylalanine, (+) or (-) phenylglycine, (+) or (-) camphanic acid or by the formation of diastereomeric carbamates by reaction of the chemical compound of the present invention with an optically active chloroformate or the like.

Additional methods for the resolving the optical isomers are known in the
25 art. Such methods include those described by *Jaques J, Collet A, & Wilen S* in "Enantiomers, Racemates, and Resolutions", John Wiley and Sons, New York (1981).

Optical active compounds can also be prepared from optical active starting materials.

30 **Methods of Producing Diazabicyclic Aryl Derivatives**

The diazabicyclic aryl derivative of the invention may be prepared by conventional methods for chemical synthesis, e.g. those described in the working examples. The starting materials for the processes described in the present application are known or may readily be prepared by conventional methods from commercially
35 available chemicals.

Also one compound of the invention can be converted to another compound of the invention using conventional methods.

The end products of the reactions described herein may be isolated by conventional techniques, e.g. by extraction, crystallisation, distillation, chromatography,
5 etc.

Biological Activity

The present invention is devoted to the provision novel ligands and modulators of the nicotinic receptors, which ligands and modulators are useful for the
10 treatment of diseases or disorders related to the cholinergic receptors, and in particular the nicotinic acetylcholine receptor (nAChR). Preferred compounds of the invention show a pronounced nicotinic acetylcholine $\alpha 7$ receptor subtype selectivity.

The compounds of the present invention may in particular be agonists, partial agonists, antagonists and/or allosteric modulators of the nicotinic acetylcholine
15 receptor.

Due to their pharmacological profile the compounds of the invention may be useful for the treatment of diseases or disorders as diverse as those related to the cholinergic system of the central nervous system (CNS), the peripheral nervous system (PNS), diseases or disorders related to smooth muscle contraction, endocrine
20 diseases or disorders, diseases or disorders related to neuro-degeneration, diseases or disorders related to inflammation, pain, and withdrawal symptoms caused by the termination of abuse of chemical substances.

The compounds of the invention may also be useful as diagnostic tools or monitoring agents in various diagnostic methods, and in particular for *in vivo* receptor
25 imaging (neuroimaging), and they may be used in labelled or unlabelled form.

In a preferred embodiment the compounds of the invention are used for the treatment of diseases, disorders, or conditions relating to the central nervous system. Such diseases or disorders includes anxiety, cognitive disorders, learning deficit, memory deficits and dysfunction, Alzheimer's disease, attention deficit, attention deficit
30 hyperactivity disorder (ADHD), Parkinson's disease, Huntington's disease, Amyotrophic Lateral Sclerosis, Gilles de la Tourette's syndrome, psychosis, depression, mania, manic depression, schizophrenia, obsessive compulsive disorders (OCD), panic disorders, eating disorders such as anorexia nervosa, bulimia and obesity, narcolepsy, nociception, AIDS-dementia, senile dementia, periferic
35 neuropathy, autism, dyslexia, tardive dyskinesia, hyperkinesia, epilepsy, bulimia, post-traumatic syndrome, social phobia, sleeping disorders, pseudodementia, Ganser's syndrome, pre-menstrual syndrome, late luteal phase syndrome, chronic fatigue syndrome, mutism, trichotillomania, and jet-lag.

In a preferred embodiment diseases, disorders, or conditions relating to the central nervous system for which the compounds of the invention are used are cognitive disorders, psychosis, schizophrenia and/or depression.

In another preferred embodiment the compounds of the invention may be
5 useful for the treatment of diseases, disorders, or conditions associated with smooth muscle contractions, including convulsive disorders, angina pectoris, premature labour, convulsions, diarrhoea, asthma, epilepsy, tardive dyskinesia, hyperkinesia, premature ejaculation, and erectile difficulty.

In yet another preferred embodiment the compounds of the invention may
10 be useful for the treatment of endocrine disorders, such as thyrotoxicosis, pheochromocytoma, hypertension and arrhythmias.

In still another preferred embodiment the compounds of the invention may be useful for the treatment of neurodegenerative disorders, including transient anoxia and induced neuro-degeneration.

15 In even another preferred embodiment the compounds of the invention may be useful for the treatment of inflammatory diseases, disorders, or conditions, including inflammatory skin disorders such as acne and rosacea, Chron's disease, inflammatory bowel disease, ulcerative colitis, and diarrhoea.

In still another preferred embodiment the compounds of the invention may
20 be useful for the treatment of mild, moderate or even severe pain of acute, chronic or recurrent character, as well as pain caused by migraine, postoperative pain, and phantom limb pain. The pain may in particular be neuropathic pain, chronic headache, central pain, pain related to diabetic neuropathy, to post therapeutic neuralgia, or to peripheral nerve injury.

25 Finally the compounds of the invention may be useful for the treatment of withdrawal symptoms caused by termination of use of addictive substances. Such addictive substances include nicotine containing products such as tobacco, opioids such as heroin, cocaine and morphine, benzodiazepines and benzodiazepine-like drugs, and alcohol. Withdrawal from addictive substances is in general a traumatic
30 experience characterised by anxiety and frustration, anger, anxiety, difficulties in concentrating, restlessness, decreased heart rate and increased appetite and weight gain.

In this context "treatment" covers treatment, prevention, prophylactics and alleviation of withdrawal symptoms and abstinence as well as treatment resulting in a
35 voluntary diminished intake of the addictive substance.

In another aspect, the compounds of the invention are used as diagnostic agents, e.g. for the identification and localisation of nicotinic receptors in various tissues.

Pharmaceutical Compositions

In another aspect the invention provides novel pharmaceutical compositions comprising a therapeutically effective amount of diazabicyclic aryl derivative of the invention.

5 While a chemical compound of the invention for use in therapy may be administered in the form of the raw chemical compound, it is preferred to introduce the active ingredient, optionally in the form of a physiologically acceptable salt, in a pharmaceutical composition together with one or more adjuvants, excipients, carriers, buffers, diluents, and/or other customary pharmaceutical auxiliaries.

10 In a preferred embodiment, the invention provides pharmaceutical compositions comprising the diazabicyclic aryl derivative of the invention, or a pharmaceutically acceptable salt or derivative thereof, together with one or more pharmaceutically acceptable carriers therefore, and, optionally, other therapeutic and/or prophylactic ingredients, known and used in the art. The carrier(s) must be
15 "acceptable" in the sense of being compatible with the other ingredients of the formulation and not harmful to the recipient thereof.

The pharmaceutical composition of the invention may be administered by any convenient route, which suits the desired therapy. Preferred routes of administration include oral administration, in particular in tablet, in capsule, in dragé, in
20 powder, or in liquid form, and parenteral administration, in particular cutaneous, subcutaneous, intramuscular, or intravenous injection. The pharmaceutical composition of the invention can be manufactured by the skilled person by use of standard methods and conventional techniques appropriate to the desired formulation. When desired, compositions adapted to give sustained release of the active ingredient
25 may be employed.

Further details on techniques for formulation and administration may be found in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing Co., Easton, PA).

The actual dosage depends on the nature and severity of the disease being
30 treated, and is within the discretion of the physician, and may be varied by titration of the dosage to the particular circumstances of this invention to produce the desired therapeutic effect. However, it is presently contemplated that pharmaceutical compositions containing of from about 0.1 to about 500 mg of active ingredient per individual dose, preferably of from about 1 to about 100 mg, most preferred of from
35 about 1 to about 10 mg, are suitable for therapeutic treatments.

The active ingredient may be administered in one or several doses per day. A satisfactory result can, in certain instances, be obtained at a dosage as low as 0.1 µg/kg i.v. and 1 µg/kg p.o. The upper limit of the dosage range is presently considered

to be about 10 mg/kg i.v. and 100 mg/kg p.o. Preferred ranges are from about 0.1 µg/kg to about 10 mg/kg/day i.v., and from about 1 µg/kg to about 100 mg/kg/day p.o.

Methods of Therapy

5 The diazabicyclic aryl derivatives of the present invention are valuable nicotinic and monoamine receptor modulators, and therefore useful for the treatment of a range of ailments involving cholinergic dysfunction as well as a range of disorders responsive to the action of nAChR modulators.

10 In another aspect the invention provides a method for the treatment, prevention or alleviation of a disease or a disorder or a condition of a living animal body, including a human, which disease, disorder or condition is responsive to modulation of cholinergic receptors and/or monoamine receptors, and which method comprises administering to such a living animal body, including a human, in need thereof an effective amount of an diazabicyclic aryl derivative of the invention.

15 In the context of this invention the term "treatment" covers treatment, prevention, prophylaxis or alleviation, and the term "disease" covers illnesses, diseases, disorders and conditions related to the disease in question.

 The preferred indications contemplated according to the invention are those stated above.

20 It is at present contemplated that suitable dosage ranges are 0.1 to 1000 milligrams daily, 10-500 milligrams daily, and especially 30-100 milligrams daily, dependent as usual upon the exact mode of administration, form in which administered, the indication toward which the administration is directed, the subject involved and the body weight of the subject involved, and further the preference and
25 experience of the physician or veterinarian in charge.

 A satisfactory result can, in certain instances, be obtained at a dosage as low as 0.005 mg/kg i.v. and 0.01 mg/kg p.o. The upper limit of the dosage range is about 10 mg/kg i.v. and 100 mg/kg p.o. Preferred ranges are from about 0.001 to about 1 mg/kg i.v. and from about 0.1 to about 10 mg/kg p.o.

30

EXAMPLES

 The invention is further illustrated with reference to the following examples, which are not intended to be in any way limiting to the scope of the invention as
35 claimed.

Example 1**Preparatory Example**

All reactions involving air sensitive reagents or intermediates were performed under nitrogen and in anhydrous solvents.

5

1,4-Diazabicyclo[3.2.2]nonan-3-one (Intermediate compound)

32.33 g (200 mmol) of 3-Quinuclidinone hydrochloride was dissolved in 75 ml of water, and to the solution of hydroxylamine hydrochloride (16.4 g; 236 mmol) and $\text{CH}_3\text{CO}_2\text{Na} \cdot 3\text{H}_2\text{O}$ (80 g; 588 mmol) was added. The mixture was stirred at 70°C for 10 hour. Then NaCl (10 g) was dissolved in the mixture and was cooled to 0°C. Separated crystals were filtered and carefully dried. The thus obtained crude 3-quinuclidone oxime (approx. 30 g) was used in the next step of the synthesis without further purification.

Polyphosphoric acid (180 g) of was heated to 100°C and crude 3-quinuclidone oxime (30 g) was added portion-wise. The reaction mixture was heated at 130°C for 20 minutes. The mixture was cooled to room temperature, and 50 ml of water was added. The mass was carefully homogenised, the mixture was poured into of ice (100 g). The mixture was made alkaline (pH 12) by adding sodium hydroxide. The mixture was extracted with chloroform (2 x 400 ml). The extract was dried over 20 sodium sulphate and the solvent was removed under reduced pressure.

Yield of the mixture of the products 1,4-diazabicyclo[3.2.2]nonan-3-one and 1,3-diazabicyclo[3.2.2]nonan-4-one was 19.02 g (68%). The mixture of isomers was crystallized from 80 ml of dioxane to yield 1,4-diazabicyclo[3.2.2]nonan-3-one (5.12 g; 18%). The solvent from filtrate was distilled off, flash chromatography (with acetone) of 25 the residue gave of 1,3-diazabicyclo[3.2.2]nonan-4-one (8.91 g; 32%).

1,4-Diazabicyclo[3.2.2]nonane [J. Med. Chem. 1993 36 2311-2320]
(Intermediate compound)

1,4-Diazabicyclo[3.2.2]nonan-3-one (5.12 g; 36 mmol) was dissolved in 30 tetrahydrofuran (50 ml), lithium aluminium hydride 2.28 g (60 mmol) was added to the solution and the reaction mixture was refluxed for 36 hours. After cooling the reaction mixture to room temperature, water (2.3 ml) was added dropwise and the mixture was filtered. The solvent was removed from the filtrate by rotavapor at reduced pressure. The formed substance was distilled with Kugelrohr (0.5 mBar; 70°C). Yield of the title 35 compound 3.11 g (68%).

Method A1-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-(2-aminophenyl)-urea free base (Compound A1)

A mixture of 1-{4-[5-(1,4-diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-(2-nitrophenyl)-urea (0.63 g; 1.32 mmol), palladium on carbon (0.60 g; 5%) and ethanol (50 ml) was stirred under hydrogen. The crude mixture was filtered and purified by silica gel chromatography, using a mixture of dichloromethane : methanol (9:1) and 1% methanol as eluent. Yield 0.50 g (85%). Mp. 174°C.

10 (1,4-Diaza-bicyclo[3.2.2]non-4-yl)-5-(4-aminophenyl)-furan-2-yl-methanone fumaric acid salt (Intermediate Compound)

The title compound was prepared according to method A from (1,4-diaza-bicyclo[3.2.2]non-4-yl)-5-(4-nitrophenyl)-furan-2-yl-methanone. Mp. 227.8°C.

15 4-Amino-N-{4-[5-(1,4-diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-benzamide free base (Compound A2)

The title compound was prepared according to Method A from N-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-4-nitro-benzamide. Mp. 151°C.

20

3-Amino-N-{4-[5-(1,4-diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-benzamide free base (Compound A3)

The title compound was prepared according to Method A from N-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-nitro-benzamide. Mp.

25 249-252°C.

(1,4-Diaza-bicyclo[3.2.2]non-4-yl)-5-(4-nitrophenyl)-furan-2-yl-methanone hydrochloric acid salt (Intermediate Compound)

A mixture of 5-(4-nitrophenyl)-2-furoyl chloride (1.0 g; 4.0 mmol), 1,4-diazabicyclo[3.2.2]nonane (0.50 g; 4.0 mmol) and 1,2-dimethoxyethane (20 ml) was stirred for 15 hours at room temperature. The title compound was filtered. Yield 1.4 g (93%). Mp. 298.2°C.

5-(4-Nitrophenyl)-2-furoyl chloride (Intermediate Compound)

35 Was prepared by stirring a mixture of 5-(4-nitrophenyl)-2-furoic acid (1.0 g; 4.3 mmol) and thionyl chloride (10 ml) at reflux for 2 hours. The mixture was evaporated and co-evaporated with anhydrous toluene. The acid chloride was used without further purification.

Method B**5-Hydroxy-1-{4-[5-(1-oxy-1,4-diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-1,5-dihydro-pyrrol-2-one (N-oxide) (Compound B1)**

A mixture of (1,4-diaza-bicyclo[3.2.2]non-4-yl)-5-(4-aminophenyl)-furan-2-yl-methanone (0.50 g; 1.6 mmol), maleic anhydride (0.24 g; 2.4 mmol) and dichloromethane (10 ml) was stirred for 4 hours at room temperature. The mixture was filtered and the title compound was isolated. Yield 0.48 g (73%). Mp. 191°C.

1-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-(2-acetylaminophenyl)-urea fumaric acid salt (Compound B2)

A mixture of 1-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-(2-aminophenyl)-urea (0.21 g; 0.47 mmol), acetic anhydride (144 mg; 1.42 mmol) and dichloromethane (20 ml) was stirred for 4 hours. Aqueous sodium hydroxide (10 ml; 1M) was added followed by extraction with dichloromethane (3 x 10 ml). The crude mixture was purified by silica gel chromatography, using a mixture of dichloromethane : methanol (9:1) and 1% methanol as eluent. Yield: 174 mg (79%). The corresponding salt was obtained by addition of a diethyl ether and methanol mixture (9:1) saturated with fumaric acid Mp. 159-169°C.

Method C**1-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-phenyl-urea free base (Compound C1)**

A mixture of (1,4-diaza-bicyclo[3.2.2]non-4-yl)-5-(4-aminophenyl)-furan-2-yl-methanone (0.50 g; 1.6 mmol), phenyl-isocyanate (498 mg; 4.18 mmol) and dichloromethane (50 ml) was stirred for 15 hours. Aqueous sodium hydroxide (10 ml; 1M) was added followed by extraction with dichloromethane (3 x 10 ml). The crude mixture was purified by silica gel chromatography, using a mixture of dichloromethane : methanol (9:1) and 1% methanol as eluent. Yield 0.16 g (23%). Mp. 153-164°C.

1-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-(2-nitrophenyl)-urea fumaric acid salt (Compound C2)

The title compound was prepared according to Method C from 2-nitrophenylisocyanate. Mp. 198-202°C.

1-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-ethyl-urea fumaric acid salt (Compound C3)

The title compound was prepared according to Method C from ethylisocyanate. Mp. 167-171°C.

1-[4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl]-3-phenylthiourea free base (Compound C4)

The title compound was prepared according to Method C from phenylisothiocyanate. Mp. 171.4-174.7°C.

5

1-[4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl]-3-(5-chloro-2-methoxy phenyl)-urea free base (Compound C5)

The title compound was prepared according to Method C from 5-chloro-2-methoxyphenylisocyanate. Isolated as an oil.

10

1-[4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl]-3-benzyl-urea free base (Compound C6)

The title compound was prepared according to Method C from benzylisocyanate. Isolated as an oil.

15

1-[4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl]-1'-benzylaminocarbonyl-3-benzyl-urea free base (Compound C7)

The title compound was prepared according to Method C from benzylisothiocyanate. Mp. 105-130°C.

20

1-[4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl]-3-(2-chlorophenyl)-urea free base (Compound C8)

The title compound was prepared according to Method C from 2-chlorophenylisocyanate. Mp. 200-211°C (decomp.).

25

1-[3-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl]-3-phenyl-urea free base (Compound C9)

The title compound was prepared according to Method C from (1,4-diaza-bicyclo[3.2.2]non-4-yl)-5-(3-aminophenyl)-furan-2-yl-methanone and phenylisocyanate. Mp. 125-130°C.

30

1-[4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl]-3-(2-fluorophenyl)-urea free base (Compound C10)

The title compound was prepared according to Method C from 2-fluorophenylisocyanate. Mp. 241°C (decomp.).

35

1-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-(3-fluorophenyl)-urea free base (Compound C11)

The title compound was prepared according to Method C from 3-fluorophenylisocyanate. Mp. 230°C.

5

1-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-(2-trifluoromethylphenyl)-urea free base (Compound C12)

The title compound was prepared according to Method C from 2-trifluoromethylphenylisocyanate. Mp. 253°C (decomp.).

10

1-[2-(3-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-ureido)-phenyl]-3-ethyl-urea fumaric acid salt (Compound C13)

The title compound was prepared according to Method C from 1-(2-amino-phenyl)-3-{4-[5-(1,4-diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-urea
15 and ethylisocyanate. Mp. 162.5-165.5°C.

1-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-(3-trifluoromethylphenyl)-urea free base (Compound C14)

The title compound was prepared according to Method C from 3-
20 trifluoromethylphenylisocyanate. Mp. 171°C.

1-[3-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl]-3-ethyl-urea fumaric acid salt (Compound C15)

The title compound was prepared according to Method C from
25 ethylisocyanate. Mp. 158-161°C.

N-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-isonicotinamide fumaric acid salt (Compound C16)

The title compound was prepared according to Method C from [5-(4-amino-
30 phenyl)-furan-2-yl]-(1,4-diaza-bicyclo[3.2.2]non-4-yl)-methanone. Mp. 250-253.5°C.

Method D

N-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-benzamide free base (Compound D1)

35

To a mixture of (1,4-diaza-bicyclo[3.2.2]non-4-yl)-5-(4-aminophenyl)-furan-2-yl-methanone (0.50 g, 1.6 mmol), diisopropylamine (415 mg, 3.2 mmol) and dichloromethane (100 ml), was added at 0°C: benzoyl chloride 0.586 g, 4.18 mmol) and stirred for 15 h at room temperature. Aqueous sodium hydroxide (10 ml; 1M) was added followed by extraction with dichloromethane (3 x 10 ml). The crude mixture was

purified by silica gel chromatography, using a mixture of dichloromethane : methanol (9:1) and 1% methanol as eluent. Yield 0.50 g (75%). Mp. 254°C.

N-{3-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-benzamide

5 fumaric acid salt (Compound D2)

The title compound was prepared according to Method D from (1,4-diaza-bicyclo[3.2.2]non-4-yl)-5-(3-aminophenyl)-furan-2-yl-methanone. Mp. 201-204°C.

N-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-2-nitro-

10 benzamide hydrochloric acid salt (Compound D3)

The title compound was prepared according to Method D from (1,4-diaza-bicyclo[3.2.2]non-4-yl)-5-(4-aminophenyl)-furan-2-yl-methanone. Mp. 283°C.

N-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-4-nitro-

15 benzamide hydrochloric acid salt (Compound D4)

The title compound was prepared according to Method D from (1,4-diaza-bicyclo[3.2.2]non-4-yl)-5-(4-aminophenyl)-furan-2-yl-methanone. Mp. 195-210°C.

N-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-nitro-

20 benzamide hydrochloric acid salt (Compound D5)

The title compound was prepared according to Method D from (1,4-diaza-bicyclo[3.2.2]non-4-yl)-5-(4-aminophenyl)-furan-2-yl-methanone. Mp. >300°C.

Method E

25 (1,4-Diaza-bicyclo[3.2.2]non-4-yl)-[5-(4-pyrrol-1-yl-phenyl)-furan-2-yl]-methanone
fumaric acid salt (Compound E1)

A mixture of 1,4-diaza-bicyclo[3.2.2]non-4-yl)-5-(4-aminophenyl)-furan-2-yl-methanone (0.50 g, 1.6 mmol), 2,5-dimethoxytetrahydrofuran (4.24 g, 64 mmol), acetic acid (0.5 ml) and dioxane (30 ml) was stirred at reflux for 20 h. Aqueous sodium
30 hydroxide (10 ml; 1M) was added followed by extraction with dichloromethane (3 x 10 ml). The crude mixture was purified by silica gel chromatography, using a mixture of dichloromethane : methanol (9:1) and 1% methanol as eluent. The corresponding salt was obtained by addition of a diethyl ether and methanol mixture (9:1) saturated with fumaric acid. Yield 0.33g (43%). Mp. 222°C.

Method F

(1,4-Diaza-bicyclo[3.2.2]non-4-yl)-[5-[4-(3-nitro-pyridin-2-ylamino)-phenyl]-furan-2-yl]-methanone fumaric acid salt (Compound F1)

A mixture of 1,4-diaza-bicyclo[3.2.2]non-4-yl-5-(4-aminophenyl)-furan-2-yl-
5 methanone (0.50 g, 1.6 mmol), 2-chloro-3-nitropyridine (0.25 g, 1.6 mmol), cesium
carbonate (0.78 g, 2.41 mmol) and NMP (0.5 ml) was stirred at 80°C for 3 days.
Aqueous sodium hydroxide (10 ml; 1M) was added followed by extraction with
dichloromethane (3 x 10 ml). The crude mixture was purified by silica gel
chromatography, using a mixture of dichloromethane : methanol (9:1) and 1%
10 methanol as eluent. Yield 110 mg (16%). The corresponding salt was obtained by
addition of a diethyl ether and methanol mixture (9:1) saturated with fumaric acid. Mp.
223°C.

Example 2

15 *In vitro* Inhibition of ³H- α -Bungarotoxine Binding in Rat Brain

In this example the affinity of the compounds of the invention for binding to
 α_7 -subtype of nicotinic receptors is determined.

α -Bungarotoxine is a peptide isolated from the venom of the Elapidae snake
Bungarus multicinctus. It has high affinity for neuronal and neuromuscular nicotinic
20 receptors, where it acts as a potent antagonist. ³H- α -Bungarotoxine labels nicotinic
acetylcholine receptors formed by the α_7 subunit isoform found in brain and the α_1
isoform in the neuromuscular junction.

Tissue preparation

25 Preparations are performed at 0-4°C. Cerebral cortices from male Wistar
rats (150-250 g) are homogenised for 10 seconds in 15 ml of 20 mM Hepes buffer
containing 118 mM NaCl, 4.8 mM KCl, 1.2 mM MgSO₄ and 2.5 mM CaCl₂ (pH 7.5)
using an Ultra-Turrax homogeniser. The tissue suspension is subjected to
centrifugation at 27,000 x g for 10 minutes. The supernatant is discarded and the pellet
30 is washed twice by centrifugation at 27,000 x g for 10 minutes in 20 ml of fresh buffer,
and the final pellet is then re-suspended in fresh buffer containing 0.01% BSA (35 ml
per g of original tissue) and used for binding assays.

Assay

35 Aliquots of 500 μ l of homogenate are added to 25 μ l of test solution and 25
 μ l of ³H- α -bungarotoxine (2 nM, final concentration) and mixed and incubated for 2
hours at 37°C. Non-specific binding is determined using (-)-nicotine (1 mM, final con-
centration). After incubation, the samples are added 5 ml of ice-cold Hepes buffer
containing 0.05% PEI and poured directly onto Whatman GF/C glass fibre filters (pre-

soaked in 0.1% PEI for at least 6 hours) under suction, and immediately washed with 2 x 5 ml ice-cold buffer.

The amount of radioactivity on the filters is determined by conventional liquid scintillation counting. Specific binding is total binding minus non-specific binding.

5 The test value is given as an IC_{50} (the concentration of the test substance which inhibits the specific binding of 3H - α -bungarotoxin by 50%).

The results of these experiments are presented in Table 1 below.


Table 1

10 Inhibition of 3H - α -Bungarotoxine Binding

Compound No.	IC_{50} (μM)
A1	0.0012
B2	0.0016
C3	0.00056

CLAIMS

1. A diazabicyclic aryl derivative represented by Formula I



5

any of its enantiomers or any mixture of its enantiomers, an N-oxide, a prodrug, or a pharmaceutically-acceptable addition salt thereof, wherein

n is 1, 2 or 3; and

10

A' and A'', independently of one another, represent an aromatic monocyclic and/or polycyclic, carbocyclic and/or heterocyclic group, optionally substituted one or more times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, carboxy, amino-carbonyl (carbamoyl), sulfamoyl and phenyl; or with another monocyclic or polycyclic, carbocyclic or heterocyclic group; which additional monocyclic or polycyclic, carbocyclic or heterocyclic group may optionally be substituted one or more times with substituents selected from the group consisting of

15

alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, carboxy, amino-carbonyl (carbamoyl), sulfamoyl and phenyl; and

25

B represents

a monocyclic heterocyclic group, optionally substituted one or more times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, oxo, carboxy, amino-carbonyl (carbamoyl), N-alkyl-amino-carbonyl (alkyl-carbamoyl), N,N-dialkyl-amino-carbonyl, alkyl-carbonyl-amino, alkyl-carbonyl-amino, sulfamoyl, phenyl or benzyl; or

30

a group of formula -NR'-B', -NR'-(C=V)-B' or -NR'-(C=V)-NR''-B'; wherein R' represents hydrogen, alkyl or a group of formula -(C=V)-NR''-B';

35

R'' represents hydrogen, alkyl, phenyl or benzyl;

V represents O, S or NR'''; wherein R''' represents hydrogen, alkyl or cyano;

and

B' represents hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, phenyl, benzyl or a monocyclic heterocyclic group; which phenyl, benzyl and heterocyclic groups are optionally substituted one or more times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, oxo, carboxy, amino-carbonyl (carbamoyl), *N*-alkyl-amino-carbonyl (alkyl-carbamoyl), *N,N*-dialkyl-amino-carbonyl, alkyl-carbonyl-amino, amino-carbonyl-amino (ureido), *N*-alkyl-amino-carbonyl-amino (*N*-alkyl-ureido), *N,N*-dialkyl-amino-carbonyl-amino (*N,N*-dialkyl-ureido), sulfamoyl, phenyl and benzyl; and

L represents

a single (covalent) bond (i.e. L is absent); or

a linking group selected from $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, $-\text{CH}=\text{CH}-$, $-\text{C}\equiv\text{C}-$, $-\text{Y}-$, $(\text{CH}_2)_m-$, $-(\text{CH}_2)_m\text{Y}-$, $-\text{CONR}''''-$, $-\text{NR}''''\text{CO}-$, $-\text{NR}''''(\text{SO}_2)-$ and $-(\text{SO}_2)\text{NR}''''-$, wherein
 Y represents $-\text{O}-$, $-\text{S}-$, $-\text{SCH}_2-$, $-\text{SO}-$, $-\text{SO}_2-$, $-\text{NR}''''-$;
 R'''' represents hydrogen or alkyl; and
 m is 0, 1, 2 or 3.

2. The diazabicyclic aryl derivative of claim 1, wherein
 n is 1, 2 or 3.

3. The diazabicyclic aryl derivative of either one of claims 1-2, wherein L represents
 a single (covalent) bond (i.e. L is absent); or
 a linking group selected from $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, $-\text{CH}=\text{CH}-$, $-\text{C}\equiv\text{C}-$, $-\text{Y}-$, $(\text{CH}_2)_m-$, $-(\text{CH}_2)_m\text{Y}-$, $-\text{CONR}''''-$, $-\text{NR}''''\text{CO}-$, $-\text{NR}''''(\text{SO}_2)-$ and $-(\text{SO}_2)\text{NR}''''-$, wherein
 Y represents $-\text{O}-$, $-\text{S}-$, $-\text{SCH}_2-$, $-\text{SO}-$, $-\text{SO}_2-$, $-\text{NR}''''-$;
 R'''' represents hydrogen or alkyl; and
 m is 0, 1, 2 or 3.

4. The diazabicyclic aryl derivative of claim 3, wherein L represents a single (covalent) bond (i.e. L is absent).

5. The diazabicyclic aryl derivative of any one of claims 1-4, wherein
 A' represents an aromatic monocyclic or polycyclic, carbocyclic or heterocyclic group, optionally substituted one or more times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-

alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, carboxy, amino-carbonyl (carbamoyl), sulfamoyl and phenyl; or with another monocyclic or polycyclic, carbocyclic or heterocyclic group; which additional monocyclic or polycyclic, carbocyclic or heterocyclic group may optionally be substituted one or more times with
5 substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, carboxy, amino-carbonyl (carbamoyl), sulfamoyl and phenyl.

10 6. The diazabicyclic aryl derivative of claim 5, wherein

A' represents an aromatic monocyclic heterocyclic group, optionally substituted one or more times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, carboxy, amino-carbonyl (carbamoyl), sulfamoyl and phenyl.

7. The diazabicyclic aryl derivative of claim 6, wherein

A' represents a furanyl, pyrrolyl, isoxazoly, 1,3,4-oxadiazoly, 1,2,3-oxadiazoly, pyridinyl, pyridinyl, pyridazinyl, indolyl, benzofuranyl, benzothienyl, quinoxalinyl or benzimidazolyl group.

8. The diazabicyclic aryl derivative of claim 7, wherein A' represents furan-2,5-diyl.

25

9. The diazabicyclic aryl derivative of any one of claims 1-8, wherein

A'' represents an aromatic monocyclic or polycyclic, carbocyclic or heterocyclic group, optionally substituted one or more times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, carboxy, amino-carbonyl (carbamoyl), sulfamoyl and phenyl; or with another monocyclic or polycyclic, carbocyclic or heterocyclic group; which additional monocyclic or polycyclic, carbocyclic or heterocyclic group may optionally be substituted one or more times with
35 substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, carboxy, amino-carbonyl (carbamoyl), sulfamoyl and phenyl.

10. The diazabicyclic aryl derivative of claim 9, wherein

A" represents a phenyl or naphthyl group; which aryl group is optionally substituted one or two times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-
5 alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, carboxy, amino-carbonyl (carbamoyl), sulfamoyl and phenyl.

11. The diazabicyclic aryl derivative of claim 10, wherein

10 A" represents a phen-1,3-diyl or phen-1,4-diyl group.

12. The diazabicyclic aryl derivative of any one of claims 1-11, wherein

B represents a monocyclic heterocyclic group, optionally substituted one or more times with substituents selected from the group consisting of alkyl, cycloalkyl,
15 cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, oxo, carboxy, amino-carbonyl (carbamoyl), *N*-alkyl-amino-carbonyl (alkyl-carbamoyl), *N,N*-dialkyl-amino-carbonyl, alkyl-carbonyl-amino, sulfamoyl, phenyl and benzyl.

20 13. The diazabicyclic aryl derivative of claim 12, wherein

B represents a monocyclic heterocyclic group selected from pyrrolidinyl, pyrrolinyl, pyrrolyl, and pyridinyl; which monocyclic heterocyclic group is optionally substituted one or two times with substituents selected from the group consisting of
25 alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, cyano, nitro, amino, oxo, carboxy, amino-carbonyl (carbamoyl), *N*-alkyl-amino-carbonyl (alkyl-carbamoyl), *N,N*-dialkyl-amino-carbonyl, alkyl-carbonyl-amino, sulfamoyl and phenyl.

30 14. The diazabicyclic aryl derivative of claim 13, wherein

B represents 3-pyrrolinyl (2,5-dihydro-pyrrolyl) or pyridinyl (pyridin-4-yl); which monocyclic heterocyclic group is optionally substituted one or two times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy, cycloalkoxy, cycloalkoxy-
35 alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, cyano, nitro, amino, oxo, carboxy, carbamoyl (amino-carbonyl), alkyl-carbamoyl (*N*-alkyl-amino-carbonyl), (*N,N*-dialkyl-amino-carbonyl), alkyl-carbonyl-amino, sulfamoyl and phenyl.

15. The diazabicyclic aryl derivative of claim 14, which is

5-Hydroxy-1-{4-[5-(1-oxy-1,4-diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-1,5-dihydro-pyrrol-2-one;

1-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-pyrrolidine-2,5-dione N-oxide; or

5 (1,4-Diaza-bicyclo[3.2.2]non-4-yl)-[5-(4-pyrrol-1-yl-phenyl)-furan-2-yl]-methanone;

or an enantiomer or a mixture of its enantiomers, or a pharmaceutically-acceptable addition salt thereof.

10 16. The diazabicyclic aryl derivative of any one of claims 1-11, wherein B represents a group of formula $-NR'-B'$, $-NR'-(C=V)-B'$ or $-NR'-(C=V)-NR''-B'$; wherein


R' represents hydrogen, alkyl or a group of formula $-(C=V)-NR''-B'$;

R'' represents hydrogen, alkyl, phenyl or benzyl;

15 V represents O, S or NR''' ; wherein R''' represents hydrogen, alkyl or cyano; and

B' represents hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, phenyl, benzyl or a monocyclic heterocyclic group; which phenyl, benzyl and heterocyclic groups are optionally substituted one or more times with substituents
20 selected from the group consisting of alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, oxo, carboxy, amino-carbonyl (carbamoyl), N-alkyl-amino-carbonyl (alkyl-carbamoyl), N,N-dialkyl-amino-carbonyl, alkyl-carbonyl-amino, amino-carbonyl-amino (ureido), N-alkyl-amino-carbonyl-amino (N-alkyl-ureido), N,N-dialkyl-amino-carbonyl-amino (N,N-dialkyl-ureido), sulfamoyl, phenyl and benzyl.

17. The diazabicyclic aryl derivative of claim 16, represented by Formula II



30

any of its enantiomers or any mixture of its enantiomers, or a prodrug, or a pharmaceutically-acceptable addition salt thereof, wherein
n, A', A'', L, R' and B' are as defined in claim 1.

35

18. The diazabicyclic aryl derivative of claim 17, wherein L represents a single (covalent) bond (i.e. L is absent);

R' represents hydrogen or alkyl; and

B' represents hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, phenyl, benzyl or a monocyclic heterocyclic group; which phenyl, benzyl and heterocyclic groups are optionally substituted one, two or three times with substituents
5 selected from the group consisting of alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, oxo, carboxy, amino-carbonyl (carbamoyl), *N*-alkyl-amino-carbonyl (alkyl-carbamoyl), *N,N*-dialkyl-amino-carbonyl, alkyl-carbonyl-amino, sulfamoyl, phenyl and benzyl.

10

19. The diazabicyclic aryl derivative of claim 18, wherein

B' represents alkyl, phenyl, benzyl, furanyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, thiadiazolyl, imidazolyl, pyrazolyl, pyridinyl, pyrimidinyl or pyridazinyl; which phenyl, benzyl and heterocyclic groups are optionally substituted one or two
15 times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, oxo, carboxy, amino-carbonyl (carbamoyl), *N*-alkyl-amino-carbonyl (alkyl-carbamoyl), *N,N*-dialkyl-amino-carbonyl, alkyl-carbonyl-amino, sulfamoyl, phenyl
20 and benzyl.

20. The diazabicyclic aryl derivative of claim 19, wherein

B' represents alkyl, phenyl, benzyl or pyridinyl; which phenyl, benzyl and pyridinyl are optionally substituted with hydroxy, alkoxy, halo, trifluoromethyl, cyano,
25 nitro, amino, *N*-alkyl-amino-carbonyl (alkyl-carbamoyl), *N,N*-dialkyl-amino-carbonyl, alkyl-carbonyl-amino, sulfamoyl, phenyl or benzyl.

21. The diazabicyclic aryl derivative of claim 17, wherein

n is 2;

30 L represents a single (covalent) bond (i.e. L is absent);

A' represents a furanyl, oxazolyl or oxadiazolyl group;

A'' represents a phenyl group; and

R' represents hydrogen or alkyl;

B' represents pyridin-2-yl, pyridin-3-yl, pyridin-4-yl; which pyridinyl may
35 optionally be substituted one or two times with alkyl, hydroxy, alkoxy, halo, trihalomethyl, trihalomethoxy, nitro and/or amino


22. The diazabicyclic aryl derivative of claim 21, which is

(1,4-Diaza-bicyclo[3.2.2]non-4-yl)-{5-[4-(3-nitro-pyridin-2-ylamino)-phenyl]-furan-2-yl}-methanone;

or an enantiomer or a mixture of its enantiomers, or a pharmaceutically-acceptable addition salt thereof.

5

23. The diazabicyclic aryl derivative of claim 16, represented by Formula III



any of its enantiomers or any mixture of its enantiomers, or a prodrug, or a
10 pharmaceutically-acceptable addition salt thereof, wherein

n , A' , A'' , L , R' , V and B' are as defined in claim 1.

24. The diazabicyclic aryl derivative of claim 23, wherein

L represents a single (covalent) bond (i.e. L is absent);

15 R' represents hydrogen or alkyl;

V represents O, S or NR''' ; wherein R''' represents hydrogen, alkyl or cyano;

and

B' represents hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, phenyl, benzyl or a monocyclic heterocyclic group; which phenyl, benzyl and
20 heterocyclic groups are optionally substituted one, two or three times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, oxo, carboxy, amino-carbonyl (carbamoyl), *N*-alkyl-amino-carbonyl (alkyl-carbamoyl), *N,N*-dialkyl-amino-
25 carbonyl, alkyl-carbonyl-amino, sulfamoyl, phenyl and benzyl.

25. The diazabicyclic aryl derivative of claim 24, wherein

B' represents phenyl, benzyl or pyridinyl; which phenyl, benzyl and pyridinyl groups are optionally substituted with halo, trifluoromethyl, cyano, nitro, amino, *N*-alkyl-
30 amino-carbonyl (alkyl-carbamoyl), *N,N*-dialkyl-amino-carbonyl, alkyl-carbonyl-amino or sulfamoyl.

26. The diazabicyclic aryl derivative of claim 25, which is

N -{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-
35 benzamide;

N-{3-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-benzamide;

N-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-2-nitro-benzamide;

5 N-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-4-nitro-benzamide;

N-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-nitro-benzamide;


4-Amino-N-{4-[5-(1,4-diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-benzamide;

3-Amino-N-{4-[5-(1,4-diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-benzamide; or

N-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-isonicotinamide;

15 or an enantiomer or a mixture of its enantiomers, or a pharmaceutically-acceptable addition salt thereof.

27. The diazabicyclic aryl derivative of claim 16, represented by Formula IV



20

any of its enantiomers or any mixture of its enantiomers, or a prodrug, or a pharmaceutically-acceptable addition salt thereof, wherein

n, A', A'', L, R', R'', V and B' are as defined in claim 1.

25

28. The diazabicyclic aryl derivative of claim 27, wherein

L represents a single (covalent) bond (i.e. L is absent);

R' represents hydrogen, alkyl or a group of formula -(C=V)-NR''-B';

R'' represents hydrogen, alkyl, phenyl or benzyl;

V represents O, S or NR'''; wherein R''' represents hydrogen, alkyl or cyano;

30 and

B' represents hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, phenyl, benzyl or a monocyclic heterocyclic group; which phenyl, benzyl and heterocyclic groups are optionally substituted one, two or three times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkyl-alkyl, hydroxy, alkoxy, hydroxyalkoxy, alkoxy-alkyl, alkoxy-alkoxy, cycloalkoxy, cycloalkoxy-alkyl, cycloalkoxy-alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, oxo, carboxy, amino-

carbonyl (carbamoyl), *N*-alkyl-amino-carbonyl (alkyl-carbamoyl), *N,N*-dialkyl-amino-carbonyl, alkyl-carbonyl-amino, amino-carbonyl-amino (ureido), *N*-alkyl-amino-carbonyl-amino (*N*-alkyl-ureido), *N,N*-dialkyl-amino-carbonyl-amino (*N,N*-dialkyl-ureido), sulfamoyl, phenyl and benzyl.

5

29. The diazabicyclic aryl derivative of claim 28, wherein B' represents alkyl, phenyl or benzyl; which phenyl and benzyl groups are optionally substituted one or two times with hydroxy, alkoxy, halo, trifluoromethyl, nitro, amino, alkyl-carbonyl-amino, amino-carbonyl-amino (ureido), *N*-alkyl-amino-carbonyl-amino (*N*-alkyl-ureido) and/or

10 *N,N*-dialkyl-amino-carbonyl-amino (*N,N*-dialkyl-ureido).

30. The diazabicyclic aryl derivative of claim 27, wherein
n is 2;

L represents a single (covalent) bond (i.e. L is absent);

15 A' represents a furanyl, oxazolyl, oxadiazolyl, thiazolyl or pyridazinyl group;

A'' represents a phenyl group; and

R' represents hydrogen, alkyl or $-(C=O)-NH-B'-$;

R'' represents hydrogen, alkyl, phenyl or benzyl;

V represents O, S or NH; and

20 B' represents a group of formula $-CH_3$, $-CH_2CH_3$, $-CH=CH_2$, $-CH=CH-CH=CH_2$, cyclopenta-1-enyl cyclopenta-2,4-dienyl, phenyl or benzyl; which phenyl and benzyl may optionally be substituted one or two times with alkyl, hydroxy, alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, amino-carbonyl (amido), *N*-alkyl-amino-carbonyl (*N*-alkyl-amido), *N,N*-dialkyl-amino-carbonyl (*N,N*-dialkyl-amido) and/or

25 alkyl-carbonyl-amino.

31. The diazabicyclic aryl derivative of claim 27, wherein
n is 2;

L represents a single (covalent) bond (i.e. L is absent);

30 A' represents a furanyl, oxazolyl, oxadiazolyl, thiazolyl or pyridazinyl group;

A'' represents a phenyl group; and

R' represents hydrogen, alkyl or $-(C=O)-NH-B'-$;

R'' represents hydrogen, alkyl, phenyl or benzyl;

V represents O, S or NH; and

35 B' represents a group of formula $-CH_3$, $-CH_2CH_3$, $-CH=CH_2$, $-CH=CH-CH=CH_2$, cyclopenta-1-enyl cyclopenta-2,4-dienyl, phenyl or benzyl; which phenyl and benzyl may optionally be substituted one or two times with alkyl, hydroxy, alkoxy, halo, trihalomethyl, trihalomethoxy, cyano, nitro, amino, amino-carbonyl (amido), *N*-alkyl-

amino-carbonyl (N-alkyl-amido), *N,N*-dialkyl-amino-carbonyl (*N,N*-dialkyl-amido) and/or alkyl-carbonyl-amino.

32. The diazabicyclic aryl derivative of claim 31, wherein

5 B' represents alkyl, phenyl, benzyl or pyridyl; which phenyl, benzyl and pyridyl groups are optionally substituted one or two times with substituents selected from the group consisting of hydroxy, alkoxy, halo, trifluoromethyl, nitro, amino, alkyl-carbonyl-amino, *N*-alkyl-amino-carbonyl-amino (*N*-alkyl-ureido), *N,N*-dialkyl-amino-carbonyl-amino (*N,N*-dialkyl-ureido) and sulfamoyl.

10

33. The diazabicyclic aryl derivative of claim 32, which is

1-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-ethyl-urea;
 1-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-phenyl-urea;
 15 1-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-(2-nitrophenyl)-urea;
 1-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-(2-acetylamino-phenyl)-urea;
 20 1-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-(2-aminophenyl)-urea;
 1-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-(5-chloro-2-methoxyphenyl)-thiourea;
 1-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-(5-chloro-2-methoxy-phenyl)-urea;
 25 1-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-benzyl-urea;
 1-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-1'-benzylaminocarbonyl-3-benzyl-urea;
 30 1-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-1'-benzylaminocarbonyl-3-benzyl-urea;
 1-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-(2-chlorophenyl)-urea;
 1-{3-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-phenyl-urea;
 35 1-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-(2-fluorophenyl)-urea;
 1-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-(3-fluorophenyl)-urea;

1-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-(2-trifluoromethylphenyl)-urea;

1-{2-[3-[4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl]-ureido]-phenyl}-3-ethyl-urea;

5 1-{4-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-(3-trifluoromethylphenyl)-urea; or

1-{3-[5-(1,4-Diaza-bicyclo[3.2.2]nonane-4-carbonyl)-furan-2-yl]-phenyl}-3-ethyl-urea;

or an enantiomer or a mixture of its enantiomers, or a pharmaceutically-
10 acceptable addition salt thereof.

34. A pharmaceutical composition comprising a therapeutically effective amount of a diazabicyclic aryl derivative of any one of claims 1-33, or a pharmaceutically-acceptable addition salt thereof, together with at least one
15 pharmaceutically-acceptable carrier or diluent.

35. Use of a diazabicyclic aryl derivative of any one of claims 1-33, or a pharmaceutically-acceptable addition salt thereof, for the manufacture of a pharmaceutical composition/medicament for the treatment, prevention or alleviation of
20 a disease or a disorder or a condition of a mammal, including a human, which disease, disorder or condition is responsive to modulation of cholinergic receptors.

36. The use according to claim 35, wherein the disease, disorder or condition relates to the central nervous system.

25 37. The use according to claim 36, wherein the disease, disorder or condition is anxiety, cognitive disorders, learning deficit, memory deficits and dysfunction, Alzheimer's disease, attention deficit, attention deficit hyperactivity disorder, Parkinson's disease, Huntington's disease, Amyotrophic Lateral Sclerosis, Gilles de la Tourette's syndrome, depression, mania, manic depression, schizophrenia, obsessive compulsive disorders (OCD), panic disorders, eating disorders such as anorexia nervosa, bulimia and obesity, narcolepsy, nociception, AIDS-dementia, senile dementia, periferic neuropathy, autism, dyslexia, tardive dyskinesia, hyperkinesia, epilepsy, bulimia, post-traumatic syndrome, social phobia, sleeping disorders,
30 pseudodementia, Ganser's syndrome, pre-menstrual syndrome, late luteal phase syndrome, chronic fatigue syndrome, mutism, trichotillomania and jet-lag.

38. The use according to claim 35, wherein the disease, disorder or condition are associated with smooth muscle contractions, including convulsive

disorders, angina pectoris, premature labour, convulsions, diarrhoea, asthma, epilepsy, tardive dyskinesia, hyperkinesia, premature ejaculation and erectile difficulty.

39. The use according to claim 35, wherein the disease, disorder or
5 condition is related to the endocrine system, such as thyrotoxicosis, pheochromocytoma, hypertension and arrhythmias.

40. The use according to claim 35, wherein the disease, disorder or
condition is a neurodegenerative disorders, including transient anoxia and induced
10 neuro-degeneration.

41. The use according to claim 35, wherein the disease, disorder or
condition is an inflammatory disorder, including inflammatory skin disorders such as
acne and rosacea, Chron's disease, inflammatory bowel disease, ulcerative colitis and
15 diarrhoea.

42. The use according to claim 35, wherein the disease, disorder or
condition is mild, moderate or even severe pain of acute, chronic or recurrent
character, as well as neuropathic pain and pain caused by migraine, postoperative
20 pain, phantom limb pain, neuropathic pain, chronic headache, central pain, pain related
to diabetic neuropathy, to post therapeutic neuralgia, or to peripheral nerve injury.

43. The use according to claim 35, wherein the disease, disorder or
condition is associated with withdrawal symptoms caused by termination of use of
25 addictive substances, including nicotine containing products such as tobacco, opioids
such as heroin, cocaine and morphine, benzodiazepines and benzodiazepine-like
drugs and alcohol.

44. A method of treatment, prevention or alleviation of a disease or a
30 disorder or a condition of a living animal body, including a human, which disorder,
disease or condition is responsive to modulation of cholinergic receptors, which
method comprises the step of administering to such a living animal body in need
thereof a therapeutically effective amount of a diazabicyclic aryl derivative of any one
of claims 1-33.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP2005/050405

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D487/08 C07D471/08 A61K31/4995 A61K31/551 A61P25/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 00/58311 A (SANOFI-SYNTHELABO; GALLET, THIERRY; JEGHAM, SAMIR; LARDENOIS, PATRICK;) 5 October 2000 (2000-10-05) cited in the application the whole document	1-44
X,P	WO 2004/016617 A (ASTRAZENECA AB; ERNST, GLEN; PHILLIPS, EIFION; SCHMIESING, RICHARD) 26 February 2004 (2004-02-26) claims ----- -/-	1-44

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

29 April 2005

Date of mailing of the international search report

18/05/2005

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Bosma, P

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP2005/050405

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>DE COSTA B R ET AL: "Synthesis and evaluation of conformationally restricted N-2-(3,4-dichlorophenyl)ethyl-N-methyl-2-(1-pyrrolidinyl)ethylamines at sigma receptors. 2. Piperazines, bicyclic amines, bridged bicyclic amines, and miscellaneous compounds"</p> <p>JOURNAL OF MEDICINAL CHEMISTRY, AMERICAN CHEMICAL SOCIETY. WASHINGTON, US, vol. 36, no. 16, 1993, pages 2311-2320, XP002125229</p> <p>ISSN: 0022-2623</p> <p>cited in the application</p> <p>the whole document</p>	1

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP2005/050405

Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claim 44 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP2005/050405

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0058311	A	05-10-2000	FR 2791678 A1	06-10-2000
			AT 232865 T	15-03-2003
			AU 3301800 A	16-10-2000
			DE 60001451 D1	27-03-2003
			DE 60001451 T2	15-01-2004
			DK 1165559 T3	10-06-2003
			EP 1165559 A1	02-01-2002
			ES 2192520 T3	16-10-2003
			WO 0058311 A1	05-10-2000
			JP 2002540208 T	26-11-2002
			SI 1165559 T1	31-08-2003
			TW 221843 B	11-10-2004
WO 2004016617	A	26-02-2004	AU 2003248592 A1	03-03-2004
			CA 2493920 A1	26-02-2004
			WO 2004016617 A1	26-02-2004

This Page Blank (usps)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

This Page Blank (uspto)